ELSEVIER

Geomorphology 27 (1999) 347–348

GEOMORPHOLOGY

Contents Volume 27, 1999

Vol. 27, Nos. 1–2

SPECIAL ISSUE

COSMOGENIC ISOTOPES IN GEMORPHOLOGY

Editorial 1

Scaling factors for the rates of production of cosmogenic nuclides for geometric shielding and attenuation at depth on sloped surfaces
J. Dunne, D. Elmore and P. Muzikar 3

Cosmogenic ^{14}C in carbonate rocks
D.A. Handwerger, T.E. Cerling and R.L. Bruhn 13

Mid-Pleistocene cosmogenic minimum-age limits for pre-Wisconsinan glacial surfaces in southwestern Minnesota and southern Baffin Island: a multiple nuclide approach
P.R. Bierman, K.A. Marsella, C. Patterson, P.T. Davis and M. Caffee 25

Dating fluvial terraces with ^{10}Be and ^{26}Al profiles: application to the Wind River, Wyoming
G.S. Hancock, R.S. Anderson, O.A. Chadwick and R.C. Finkel 41

Use of cosmogenic ^{35}S for comparing ages of water from three alpine–subalpine basins in the Colorado Front Range
J.K. Sueker, J.T. Turk and R.L. Michel 61

Determining the times and distances of particle transit in a mountain stream using fallout radionuclides
E.C. Bonniwell, G. Matisoff and P.J. Whiting 75

Cosmogenic ^{3}He ages and frequency of late Holocene debris flows from Prospect Canyon, Grand Canyon, USA
T.E. Cerling, R.H. Webb, R.J. Poreda, A.D. Rigby and T.S. Melis 93

Long-term rates of denudation in the Dry Valleys, Transantarctic Mountains, southern Victoria Land, Antarctica based on in-situ-produced cosmogenic ^{21}Ne
M.A. Summerfield, F.M. Stuart, H.A.P. Cockburn, D.E. Sugden, G.H. Denton, T. Dunai and D.R. Marchant 113

Estimates of the rate of regolith production using ^{10}Be and ^{26}Al from an alpine hillslope
E.E. Small, R.S. Anderson and G.S. Hancock 131

Cosmogenic nuclides, topography, and the spatial variation of soil depth
A.M. Heimsath, W.E. Dietrich, K. Nishiizumi and R.C. Finkel 151

Vol. 27, Nos. 3–4

Research Papers

A new approach to the relief of Great Britain. III. Derivation of the contribution of neotectonic movements and exceptional regional denudation to the present relief
K. Clayton and N. Shamoon 173

The morphologic structure of step–pools in mountain streams
A. Chin 191

Rates of stream incision in the middle part of the Arkansas River basin based on late Tertiary to mid-Pleistocene volcanic ash
P.A. Ward III and B.J. Carter 205

Velocity reversals and sediment sorting in pools and riffles controlled by channel constrictions
D.M. Thompson, E.E. Wohl and R.D. Jarrett 229

Active tectonics of the Loreto area, Baja California Sur, Mexico
L. Mayer and K.R. Vincent 243

The geomorphic evolution of slopes and sediment chutes on forereefs
D.B. Hall 257
Observations regarding the movement of barchan sand dunes in the Nazca to Tanaca area of southern Peru
S. Parker Gay Jr. 279
The Alemania rockfall dam: A record of a mid-holocene earthquake and catastrophic flood in northwestern Argentina
W.J. Wayne. 295
Lacustrine sedimentation in a monsoon environment: the record from Phewa Tal, middle mountain region of Nepal
J. Ross and R. Gilbert 307
The gravel–sand transition in a disturbed catchment
A.D. Knighton 325

Newsletter of the IAG 343

Contents Volume 27, 1999 347